



Maximizing products of linear forms, and the permanent of positive semidefinite matrices

Chenyang Yuan¹ · Pablo A. Parrilo¹

Received: 11 April 2020 / Accepted: 4 January 2021

© Springer-Verlag GmbH Germany, part of Springer Nature and Mathematical Optimization Society 2021

Abstract

We study the convex relaxation of a polynomial optimization problem, maximizing a product of linear forms over the complex sphere. We show that this convex program is also a relaxation of the permanent of Hermitian positive semidefinite (HPSD) matrices. By analyzing a constructive randomized rounding algorithm, we obtain an improved multiplicative approximation factor to the permanent of HPSD matrices, as well as computationally efficient certificates for this approximation. We also propose an analog of van der Waerden’s conjecture for HPSD matrices, where the polynomial optimization problem is interpreted as a relaxation of the permanent.

1 Introduction

We study the problem of maximizing a product of linear forms on the complex $(n - 1)$ -sphere of radius \sqrt{n} :

$$r(A) \equiv \max_{\|x\|^2=n} \prod_{i=1}^n |\langle x, v_i \rangle|^2, \quad (1)$$

where $A = V^\dagger V$ and v_i are the columns of V . We show that the natural convex relaxation of (1),

$$\max \prod_{i=1}^n v_i^\dagger P v_i \text{ s.t. } \text{Tr}(P) = n, P \succeq 0, \quad (2)$$

✉ Chenyang Yuan
ycy@mit.edu

Pablo A. Parrilo
parrilo@mit.edu

¹ Massachusetts Institute of Technology, Cambridge, MA, USA

is also a relaxation of the permanent of A , which is defined by

$$\text{per}(A) = \sum_{\sigma \in \mathcal{S}_n} \prod_{i=1}^n A_{i, \sigma(i)}, \quad (3)$$

where the sum is over all $n!$ permutations of n elements. Computing the permanent exactly is #P-hard [14], and approximation efforts have been focused on classes of matrices with computationally efficient certificates of permanent non-negativity. For matrices with non-negative entries, [9] gave a randomized algorithm achieving a $(1 + \epsilon)$ -approximation. There has been recent interest in approximating the permanent of HPSD matrices due to their applications in quantum information [7]. The work by [1] gave the first polynomial-time algorithm for approximating the permanent of HPSD matrices with a simply exponential multiplicative approximation factor of $\frac{n!}{n^n} e^{-n\gamma}$, where $\gamma \approx 0.577$ is the Euler–Mascheroni constant. Their algorithm is based on the following convex program relaxation of the permanent.

Definition 1.1 Given a HPSD matrix $A \in \mathbb{C}^{n \times n}$, we define $\text{rel}(A)$ as the solution to the optimization problem:

$$\text{rel}(A) \equiv \begin{cases} \min \prod_{i=1}^n D_{ii} \\ \text{s.t. } A \preceq D, \quad D \text{ is diagonal} \end{cases} \quad (4)$$

In this paper, we show that $\text{rel}(A)$ is equivalent to the convex relaxation (2). Our main result, Theorem 4.4, uses this connection between the polynomial optimization problem (1) and $\text{per}(A)$ to provide a new analysis of the approximation of $\text{per}(A)$ in terms of the rank of the optimal solution to (2). By bounding this rank, we prove an improved approximation factor for all finite n :

Corollary 1.2 Given a HPSD matrix $A \in \mathbb{C}^{n \times n}$, $\text{rel}(A)$ is an $\frac{n!}{n^n} e^{-nL_r}$ -approximation to $\text{per}(A)$:

$$\frac{n!}{n^n} e^{-nL_r} \text{rel}(A) \leq \text{per}(A) \leq \text{rel}(A)$$

where $r = O(\sqrt{n})$, $L_r = H_{r-1} - \log(r)$, and $H_r = \sum_{k=1}^r \frac{1}{k}$ is the r -th harmonic number.

From the definition of the Euler–Mascheroni constant, $\lim_{n \rightarrow \infty} L_r = \gamma$. For any finite n , $L_r < \gamma$ and thus $\frac{n!}{n^n} e^{-nL_r} > \frac{n!}{n^n} e^{-n\gamma}$. More precisely, using Proposition A.1, we can show that this is a $e^{O(\sqrt{n})}$ multiplicative improvement. [1] also constructed a series of matrices A_k such that $(\text{rel}(A_k)/\text{per}(A_k))^{1/n} \rightarrow e^{1+\gamma}$ as $k \rightarrow \infty$. However since this result only rules out improvements on the order of $e^{O(n)}$, it does not contradict Corollary 1.2.

In Sect. 3 we analyze the convex relaxation of (1), describe a rounding procedure and prove its approximation factor. In particular we prove that

$$e^{-nL_r} \text{rel}(A) \leq r(A) \leq \text{rel}(A).$$

In Sect. 4 we prove Theorem 4.4. We first show that the convex relaxation of $r(A)$ is equivalent to $\text{rel}(A)$. Then using the vector produced by the rounding procedure of the relaxation, we construct a rank-1 matrix whose permanent lower bounds $\text{per}(A)$, thus showing that $\text{rel}(A)$ also well-approximates $\text{per}(A)$. Note that in [1] only the existence of this rank-1 matrix is shown, but in our analysis we provide an explicit construction of a rank-1 matrix whose permanent lower bounds $\text{per}(A)$. This combined with the diagonal matrix in (4) whose permanent upper-bounds $\text{per}(A)$ certifies the approximation. In Sect. 5 we explore reasons why the convex relaxation of (1) is equivalent to $\text{rel}(A)$. We conjecture that (1) is itself a $\frac{n!}{n^n}$ approximation to $\text{per}(A)$, explain why it is an analogue of van der Waerden’s conjecture, and show that it is implied by another long-standing permanent conjecture.

2 Preliminaries

For any $x \in \mathbb{C}$, let x^* be its complex conjugate, and $|x|^2 = xx^*$. For any matrix $A \in \mathbb{C}^{n \times m}$, let $A^\dagger = (A^*)^T$ be its conjugate transpose. Given $a, b \in \mathbb{C}^n$, let $\langle a, b \rangle = a^\dagger b$ be the inner product on the Hilbert space \mathbb{C}^n , and $\|a\|^2 = \langle a, a \rangle$. Let $S_{\mathbb{C}}(n) = \{x \in \mathbb{C}^n \mid \|x\|^2 = n\}$ be the complex sphere in n dimensions of radius \sqrt{n} . A matrix A is Hermitian if $A = A^\dagger$, and is Hermitian positive semidefinite (HPSD) if in addition $x^\dagger Ax \geq 0$ for all $x \in \mathbb{C}^n$. We can also denote this as $A \succeq 0$. The \succeq operator induces a partial order called the Löwner order, where $A \succeq B$ if $A - B \succeq 0$. If $A \succeq 0$, it can be factorized as $A = L^\dagger L$, where $L \in \mathbb{C}^{n \times n}$ (for example by the Cholesky decomposition).

2.1 Circularly-symmetric gaussian random variables

In this paper we will use a few results involving vectors of circularly-symmetric valued Gaussian variables.

Definition 2.1 (Circularly-symmetric Gaussian random vector) The complex-valued Gaussian random variable $Z = Z_r + iZ_c$ is circularly-symmetric if Z_r and Z_c are i.i.d. drawn from $\mathcal{N}(0, \frac{1}{2})$. The random vector $\mathbf{Z} = [Z_1, \dots, Z_n]^T$ is drawn from the distribution $\mathcal{CN}(0, \Sigma)$ if Z_i are i.i.d. circularly-symmetric Gaussians and $\mathbb{E}[\mathbf{Z}\mathbf{Z}^\dagger] = \Sigma$.

The name circularly-symmetric comes from the fact that \mathbf{Z} is invariant under rotations in the complex plane, meaning that $e^{i\theta}\mathbf{Z}$ has the same distribution as \mathbf{Z} for all real θ . All complex multivariate Gaussians in this paper are circularly symmetric. Similar to real multivariate Gaussians, a linear transform on the random vector induces a congruence transform on the covariance matrix.

Proposition 2.2 (Linear transformations of complex multivariate Gaussians) *Given $\mathbf{Z} \sim \mathcal{CN}(0, \Sigma)$ and any complex matrix A , $A\mathbf{Z}$ is also circularly symmetric and has the distribution $\mathcal{CN}(0, A\Sigma A^\dagger)$.*

The proof of this proposition and more about complex multivariate Gaussians can be found in [6]. In particular, this tells us that $\mathbf{Z} \sim \mathcal{CN}(0, I)$ is invariant under unitary transformations.

In the analysis of our rounding procedure, we use some results about the gamma distribution.

Fact 2.3 (Expectation of log of gamma random variable) *Let $X \sim \text{Gamma}(\alpha, \beta)$ be drawn from the gamma distribution, with density $p(x; \alpha, \beta) = \Gamma(\alpha)^{-1} \beta^\alpha x^{\alpha-1} e^{-\beta x}$. Then*

$$\mathbb{E}[\log X] = \psi(\alpha) - \log(\beta),$$

where $\psi(x) = \frac{d}{dx} \log \Gamma(x)$ is the digamma function.

This follows from the fact that the gamma distribution is an exponential family, and $\log x$ is a sufficient statistic (see section 2.2 of [10] for more details). Next we prove a useful identity.

Fact 2.4 *Let $[z_1, \dots, z_r]^T \sim \mathcal{CN}(0, I_r)$, $H_n = \sum_{k=1}^n \frac{1}{k}$ be the n -th harmonic number and $\gamma = \lim_{n \rightarrow \infty} (H_n - \log n)$ be the Euler-Mascheroni constant. Then*

$$\mathbb{E} \log \left(\frac{1}{r} \sum_{i=1}^r |z_i|^2 \right) = H_{r-1} - \gamma - \log(r) = L_r - \gamma.$$

Proof $\sum_{i=1}^r 2|z_i|^2$ is distributed as a chi-squared distribution with $2r$ degrees of freedom, which is equivalent to $\text{Gamma}(r, \frac{1}{2})$. Using Fact 2.3, $\mathbb{E} \log(\sum_{i=1}^r |z_i|^2) = \psi(r)$. Since $\psi(1) = -\gamma$ by Gauss's digamma theorem, the recurrence relation of the gamma function shows that for all positive integers r , $\psi(r) = H_{r-1} - \gamma$. \square

Integrating a homogeneous polynomial over the complex sphere is equivalent to taking its expectation with respect to $x \sim \mathcal{CN}(0, I)$, up to a correction factor. This factor can be found by computing moments of a chi-squared distribution.

Fact 2.5 *Let $p(x)$ be a degree d homogeneous polynomial in n variables, $\mu_n(x)$ be the measure associated with the random variable $x \sim \mathcal{CN}(0, I_n)$. Then*

$$\int_{\mathbb{C}^n} |p(x)|^2 d\mu_n(x) = \frac{(n+d-1)!}{n^n(n-1)!} \int_{S_{\mathbb{C}}(n)} |p(x)|^2 dx.$$

2.2 Permanent of HPSD matrices

One remarkable property of the permanent of HPSD matrices is that it respects the Löwner order. See section 2.3 of [1] for a proof.

Proposition 2.6 *If $A \geq B \geq 0$, then $\text{per}(A) \geq \text{per}(B) \geq 0$.*

We can efficiently compute the permanent of rank-1 matrices. The following proposition immediately follows from the definition of the permanent in (3).

Proposition 2.7 For any $v \in \mathbb{C}^n$, $\text{per}(vv^\dagger) = n! \prod_{i=1}^n |v_i|^2$.

The permanent of HPSD matrices also has an integral representation using complex multivariate Gaussians. See section 4 of [3] for more details and a proof.

Proposition 2.8 Let $\mu_n(x)$ be the measure associated with the random variable $x \sim \mathcal{CN}(0, I_n)$, and $S_{\mathbb{C}}(n)$ be the complex $(n - 1)$ -sphere with radius \sqrt{n} . For any HPSD $A = V^\dagger V$, where v_i are the columns of V ,

$$\text{per}(A) = \int_{\mathbb{C}^n} \prod_{i=1}^n |\langle v_i, x \rangle|^2 d\mu_n(x) = \frac{(2n - 1)!}{n^n(n - 1)!} \int_{S_{\mathbb{C}}(n)} \prod_{i=1}^n |\langle v_i, x \rangle|^2 dx.$$

3 Convex relaxation and rounding

In this section we analyze the convex relaxation (2) and a natural rounding algorithm for maximizing a product of linear forms over the complex sphere.

Remark 3.1 Both $r(A)$ and $\text{rel}(A)$ are independent of the factorization of $A = V^\dagger V$. This is because any two different factorizations of $A = V_1^\dagger V_1 = V_2^\dagger V_2$ are related by a unitary transform $V_1 = UV_2$ for some unitary matrix U ¹. This induces a change of variables $x \mapsto U^\dagger x$ in (1) but does not change the value of $r(A)$.

Lemma 3.2 Any $A \geq 0$ can be factorized as $A = V^\dagger V$, where v_i are the columns of V . Consider the following pair of convex programs:

$$\mu^*(A) \equiv \min \lambda^n \text{ s.t. } \begin{cases} V \text{Diag}(\alpha)V^\dagger \leq \lambda I_n \\ \prod_{i=1}^n \alpha_i \geq 1 \\ \alpha_i > 0 \end{cases} \tag{5}$$

$$\nu^*(A) \equiv \max \prod_{i=1}^n v_i^\dagger P v_i \text{ s.t. } \begin{cases} \text{Tr}(P) = n \\ P^\dagger = P \\ P \geq 0 \end{cases} \tag{6}$$

Then $r(A) \leq \nu^*(A) = \mu^*(A)$, thus the convex programs are relaxations of $r(A)$ (see Eq. (1)).

Proof If we add a rank-1 constraint to (6), we get (1), showing that $r(A) \leq \nu^*(A)$. Suppose we have feasible solutions λ, α_i and P to (5) and (6) respectively. Then

$$\prod_{i=1}^n v_i^\dagger P v_i = \prod_{i=1}^n \alpha_i v_i^\dagger P v_i \leq \left(\frac{1}{n} \sum_{i=1}^n \alpha_i v_i^\dagger P v_i \right)^n \leq \left(\frac{\text{Tr}(P)\lambda}{n} \right)^n = \lambda^n,$$

showing weak duality, i.e. $\nu^*(A) \leq \mu^*(A)$. Since (6) comes from taking the dual of (5) and has a strictly feasible solution, strong duality holds, i.e. $\nu^*(A) = \mu^*(A)$. If $P = xx^\dagger$ is rank-1, then $v_i^\dagger P v_i = |\langle x, v_i \rangle|^2$, thus in (6) the variable P can be interpreted as the convex relaxation of the rank-1 constraint in (1). \square

¹ We are assuming here that $V_1, V_2 \in \mathbb{C}^{n \times n}$ even if $\text{rank}(A) < n$, padding with zero columns if necessary.

Although (5) and (6) have non-linear objective functions and are not semidefinite programs in standard form, the geometric mean constraint/objective in them can be converted to second-order conic constraints after a change of variables [11]. They can also be solved efficiently with convex programming techniques such as interior point methods (see [15]). Our main result (Theorem 4.4) is proven with the following analysis of a randomized rounding procedure to the convex relaxation of the product of linear forms. This produces a vector that gives an e^{-nL_r} -approximation to (1).

Theorem 3.3 *Given a matrix $A \succeq 0$, let $v^*(A)$ be the optimum of (6), with optimum achieved by $P^* = UU^\dagger$. Suppose P^* has rank r , therefore $U \in \mathbb{C}^{n \times r}$. If we produce a vector $y \in S_{\mathbb{C}}(n)$ using the following procedure:*

1. Sample $z \in \mathbb{C}^r$ uniformly at random from the complex multivariate Gaussian $\mathcal{CN}(0, I_r)$
2. Return the normalized vector $y = \sqrt{n}Uz / \|Uz\|$

Recalling that $L_r = H_{r-1} - \log r$, we have the following lower bound on the expected value of the objective:

$$\mathbb{E}_z \left[\prod_{i=1}^n |\langle v_i, y \rangle|^2 \right] \geq e^{-nL_r} v^*(A)$$

Proof We use Jensen’s inequality to bound the expectation:

$$\begin{aligned} \mathbb{E}_z \left[\prod_{i=1}^n \frac{n |\langle v_i, Uz \rangle|^2}{\|Uz\|^2} \right] &= \mathbb{E}_z \left[\exp \left(\sum_{i=1}^n (\log |\langle v_i, Uz \rangle|^2 - \log z^\dagger U^\dagger Uz + \log n) \right) \right] \\ &\geq \exp \left(\sum_{i=1}^n (\mathbb{E}_z \log |\langle v_i, Uz \rangle|^2 - \mathbb{E}_z \log z^\dagger U^\dagger Uz + \log n) \right) \end{aligned}$$

We can exactly compute the first expectation:

$$\begin{aligned} \mathbb{E}_z \log |\langle v_i, Uz \rangle|^2 &= \log v_i^\dagger U U^\dagger v_i + \mathbb{E}_z \log \left| \left\langle U^\dagger v_i / \|U^\dagger v_i\|, z \right\rangle \right|^2 \\ &= \log v_i^\dagger U U^\dagger v_i + \mathbb{E}_z \log |z_1|^2 \\ &= \log v_i^\dagger P^* v_i - \gamma \end{aligned}$$

Where the first equality follows from normalizing $U^\dagger v_i$, the second equality follows from the rotational symmetry of the complex multivariate Gaussian since $U^\dagger v_i / \|U^\dagger v_i\|$ is a unit vector, and the third equality follows from Fact 2.4 for $r = 1$. Let $\lambda_1, \dots, \lambda_r$ be the eigenvalues of $U^\dagger U$. Then

$$\mathbb{E}_z \log z^\dagger U^\dagger Uz = \mathbb{E}_z \log \left(\sum_{i=1}^r \lambda_i |z_i|^2 \right) \leq \mathbb{E}_z \log \left(\frac{n}{r} \sum_{i=1}^r |z_i|^2 \right) = H_{r-1} - \gamma + \log \left(\frac{n}{r} \right),$$

where the first equality follows from the invariance of the complex multivariate Gaussian under unitary transformations (see Proposition 2.2), and the second equality follows from Fact 2.4. Next we prove the inequality. Since $\text{Tr}(U^\dagger U) = \text{Tr}(P^*) = n$, $\lambda = (\lambda_1, \dots, \lambda_r)$ lies on the scaled r -simplex. The function $\lambda \mapsto \mathbb{E}_z \log \left(\sum_{i=1}^r \lambda_i |z_i|^2 \right)$ is concave on the scaled r -simplex and is symmetric with respect to all permutations of the coordinates of λ , therefore it is maximized when all $\lambda_i = \frac{n}{r}$. Finally we put the above together, along with the fact that $\prod_{i=1}^n v_i^\dagger P^* v_i = v^*(A)$, to prove the theorem. \square

4 Approximating the permanent

We present a new analysis of the relaxation of the permanent of HPSD matrices in [1]. First we show that $\text{rel}(A)$ is a relaxation of $\text{per}(A)$.

Lemma 4.1 *Given any $A \geq 0$,*

$$\text{per}(A) \leq \text{rel}(A).$$

Proof Using the monotonicity of the permanent with respect to the Löwner order (Proposition 2.6), $A \preceq D$ implies that $\text{per}(A) \leq \text{per}(D)$. Since D is diagonal, $\text{per}(D) = \prod_i D_{ii}$, showing that the permanent is always bounded by $\text{rel}(A)$. \square

Next we show that $\text{rel}(A)$ is equivalent to the convex relaxation of (1).

Lemma 4.2 *Recall that $\mu^*(A) = v^*(A)$ is the optimal value of the convex relaxation in Lemma 3.2. Then*

$$\text{rel}(A) = \mu^*(A) = v^*(A).$$

Proof By a scaling argument, the optimum of (5) is achieved when $\prod_i \alpha_i = 1$. Taking Schur complements, $V \text{Diag}(\alpha) V^\dagger \preceq \lambda I_n$ is equivalent to $\lambda \text{Diag}(\alpha)^{-1} \succeq V^\dagger V = A$. Thus by making the substitution $D_{ii} = \lambda/\alpha_i$ and noting that $\prod_i D_{ii} = \lambda^n$, we show that $\text{rel}(A) = \mu^*(A)$. \square

The following lemma shows that given any vector $y \in S_{\mathbb{C}}(n)$ returned by the rounding algorithm, we can construct a lower bound on $\text{per}(A)$.

Lemma 4.3 *Given HPSD $A = V^\dagger V \in \mathbb{C}^{n \times n}$, where v_i are columns of V , and a vector $y \in S_{\mathbb{C}}(n)$,*

$$\frac{n!}{n^n} \prod_{i=1}^n |\langle v_i, y \rangle|^2 \leq \text{per}(A).$$

Proof Since $\|y\|^2 = n$, $yy^\dagger \preceq nI$ and $V^\dagger yy^\dagger V \preceq nV^\dagger V$. Thus $n^{-n} \text{per}(V^\dagger yy^\dagger V) \leq \text{per}(V^\dagger V) = \text{per}(A)$. Since $V^\dagger yy^\dagger V$ is a rank-1 matrix, its permanent is $n! \prod_i |\langle v_i, y \rangle|^2$ by Proposition 2.7. \square

Now we can state our result about approximating the permanent of a HPSD matrix.

Theorem 4.4 *Given a HPSD matrix $A \in \mathbb{C}^{n \times n}$, $\text{rel}(A)$ is a relaxation of $\text{per}(A)$ computable in polynomial-time by convex programs (5) or (6). Let r be the rank of P^* , the solution to (6). Then $\text{rel}(A)$ is an $\frac{n!}{n^n} e^{-nL_r}$ -approximation to $\text{per}(A)$:*

$$\frac{n!}{n^n} e^{-nL_r} \text{rel}(A) \leq \text{per}(A) \leq \text{rel}(A) \quad (7)$$

Next we state a result that we will use to bound the rank of P^* .

Lemma 4.5 (Theorem 2.2 in [2]) *Suppose there is a non-zero solution X to the system of equations $\{X \geq 0, \text{Tr}(A_i X) = b_i, i = 1, \dots, d\}$, where A_i is Hermitian and $b_i \in \mathbb{C}$. If $d < (r + 1)^2$, then one can find in polynomial time another solution X' where $\text{rank}(X') = r$.*

We can now prove Corollary 1.2.

Proof of Corollary 1.2 Given a solution P to (6), any HPSD matrix P' that satisfy the $n + 1$ equalities $\text{Tr}(v_i v_i^\dagger P') = v_i^\dagger P v_i$ and $\text{Tr}(P') = n$ will have the same objective value as that of P . Applying Lemma 4.5, we can find in polynomial time an optimal solution P^* with $\text{rank}(P^*) \leq O(\sqrt{n})$. We then apply Theorem 4.4. \square

Finally we prove Theorem 4.4.

Proof of Theorem 4.4 We use the vector y produced in the rounding procedure in Theorem 3.3 to construct a rank-1 matrix $V^\dagger y y^\dagger V$. We then compare the permanent of this matrix to $\text{per}(A)$ and $\text{rel}(A)$:

$$\frac{n!}{n^n} e^{-nL_r} \text{rel}(A) \stackrel{1}{=} \frac{n!}{n^n} e^{-nL_r} v^*(A) \stackrel{2}{\leq} \frac{n!}{n^n} \mathbb{E} \left[\prod_{i=1}^n |\langle v_i, y \rangle|^2 \right] \stackrel{3}{\leq} \text{per}(A) \stackrel{4}{\leq} \text{rel}(A)$$

1. Apply Lemma 4.2.
2. Apply Theorem 3.3.
3. Lemma 4.3 shows that for any vector $y \in S_{\mathbb{C}}(n)$, $\text{per}(A) \geq \frac{n!}{n^n} \prod_{i=1}^n |\langle v_i, y \rangle|^2$. This is also true when taking an expectation of any distribution supported on $S_{\mathbb{C}}(n)$.
4. Apply Lemma 4.1.

\square

4.1 Low rank instances

There are structured classes of HPSD matrices where we can prove a priori that the rank of P^* is low and thus a better approximation ratio can be obtained. For example, it is easy to show that $\text{rank}(P^*) \leq \text{rank}(A)$. Often such instances also have additional symmetry, such as the class of circulant matrices.




Fig. 1 Plot of average of $\text{rank}(P^*)$ (along with error bars indicating standard deviation) of 50 random instances as a function of n . $A = V^\dagger V$ is constructed by sampling each entry of $V \in \mathbb{R}^{n \times n}$ from a standard Gaussian. The theoretical upper bound for $\text{rank}(P^*)$ in Lemma 4.5 is also shown for comparison

Corollary 4.6 *A square matrix is circulant if each row is cyclically shifted one position to the right compared to the previous row. If $A \in \mathbb{C}^{n \times n}$ is HPSD and circulant, then there is a solution P^* to (6) where $\text{rank}(P^*) = 1$ and we have the bound*

$$\frac{n!}{n^n} \text{rel}(A) \leq \text{per}(A) \leq \text{rel}(A).$$

Proof Since $A = V^\dagger V$ is circulant it is invariant under the map $A_{i,j} \mapsto A_{(i+1 \bmod n), (j+1 \bmod n)}$. Suppose we have an optimal solution D^* to $\text{rel}(A)$ in (4), where D^* is a diagonal matrix satisfying $A \leq D^*$. We then average over all cyclic shifts of D^* to show that $D = \lambda I$ is also optimal, which corresponds to $\alpha_i = 1$ in (5), with an optimal solution P of (6) satisfying the complementary slackness condition of $\text{Tr}(P V V^\dagger) = n\lambda$. This shows that $P = v v^\dagger$ is also a solution, where v is a suitable multiple of the top eigenvector of $V V^\dagger$. \square

We also observed experimentally that $\text{rank}(P^*)$ is small for random A . Figure 1 plots this rank as a function of n , for instances of A drawn from the Gaussian orthogonal ensemble. The results suggest that $\text{rank}(P^*)$ for these random instances grows slower than $O(\sqrt{n})$.

5 A conjecture

Our analysis of $\text{rel}(A)$ was inspired by the optimization problem (1), maximizing a product of linear forms over the complex sphere. We conjecture that the exact solution to this optimization problem is a tighter relaxation of the permanent.

Conjecture 5.1 *Given $A = V^\dagger V$, where v_i are the columns of V , recall that $r(A)$ is the maximum of a product of linear forms as defined in (1). Then*

$$\frac{n!}{n^n} r(A) \leq \text{per}(A) \leq r(A). \tag{8}$$

If the matrix A is scaled so that $r(A) = 1$, then (8) is exactly the same bounds given by the van der Waerden's conjecture for doubly stochastic matrices (proved by [4,5,8]). The lower bound follows from Lemma 4.3, but the upper bound cannot be proven by naively applying Proposition 2.8 and bounding the integral over the complex sphere by its maximum. However, we can show that the upper bound is implied by another conjecture on permanents:

Conjecture 5.2 (Pate's conjecture [12]) *Given any $n \times n$ HPSD matrix A , let $A \otimes J_k$ be the Kronecker product of A with the $k \times k$ all-ones matrix. Then*

$$\text{per}(A \otimes J_k) \geq \text{per}(A)^k (k!)^n. \quad (9)$$

This conjecture has been proved in the case where $n = 2$, see [16] for a survey of subsequent progress on this conjecture. Using the integral representation of the permanent (Proposition 2.8), we can write (9) as:

$$\mathbb{E}_{x \sim \mathcal{CN}(0, I_n)} \left[\prod_{i=1}^n |\langle v_i, x \rangle|^{2k} \right]^{1/k} \geq \text{per}(A) \mathbb{E}_{x \sim \mathcal{CN}(0, I_n)} \left[\prod_{i=1}^n |x_i|^{2k} \right]^{1/k}$$

Since both expectations are taken over homogeneous polynomials of degree d , we can apply Fact 2.5, take $k \rightarrow \infty$ and get:

$$\max_{\|x\|^2=n} \prod_{i=1}^n |\langle v_i, x \rangle|^2 \geq \text{per}(A) \max_{\|x\|^2=n} \prod_{i=1}^n |x_i|^2 = \text{per}(A).$$

6 Discussion and conclusion

There are a few interesting directions that stem from this work. For random A (i.e. drawn from the Gaussian orthogonal ensemble), numerical experiments in Sect. 4.1 suggest that $\text{rank}(P^*)$ is very small compared to \sqrt{n} . It would be interesting to provide concrete bounds on the rank of random instances. One might also ask if we can construct sequences of matrices A_k of increasing size but with fixed rank r , where $(\text{rel}(A_k)/\text{per}(A_k))^{1/n} \rightarrow e^{1+Lr}$. This is related to the question called the *linear polarization constant of Hilbert spaces*, see [13] for such a construction and its analysis.

The main result of this paper uses the connection between the permanent and the optimization of a product of linear forms over the sphere (1). Although it is natural to conjecture the hardness of computing $r(A)$, we do not know of any formal results establishing this. We also proposed Conjecture 5.1 which would explain why this optimization problem is intimately related to the permanent. Better understanding of this problem may lead to further insights about the permanent of HPSD matrices.

Acknowledgements This work was partially supported by NSF AF 1565235.




Fig. 2 Illustration of the asymptotics of L_r . Δ_r is the area between the curve $y = \frac{1}{x}$ and the rectangle of height $\frac{1}{r+1}$. The lower dotted lines are tangent to the curve at $r + 1$ and $r + 2$ respectively

Appendix: Asymptotics of the approximation factor

Proposition A.1 For all positive integers r ,

$$\frac{1}{2r} < \gamma - L_r < \frac{r + 2}{2r(r + 1)}. \tag{10}$$

Proof It is easy to see that (10) follows from

$$\frac{1}{2(r + 1)} < H_n - \log(r) - \gamma < \frac{1}{2r}.$$

From Fig. 2, we can see that $H_n - \log(r) - \gamma = \sum_{k=r}^{\infty} \Delta_k$. The upper bound is given by computing the sum of the areas of the larger triangles:

$$\sum_{k=r}^{\infty} \Delta_k < \sum_{k=r}^{\infty} \frac{1}{2} \left(\frac{1}{k} - \frac{1}{k + 1} \right) = \sum_{k=r}^{\infty} \frac{1}{2k(k + 1)} = \frac{1}{2r}$$

The lower bound is given by computing the sum of the areas of the smaller triangles:

$$\sum_{k=r}^{\infty} \Delta_k > \sum_{k=r}^{\infty} \frac{1}{2} \frac{1}{(k + 1)^2} > \sum_{k=r}^{\infty} \frac{1}{2(k + 1)(k + 2)} = \frac{1}{2(r + 1)}$$

□

References

1. Ai, W., Huang, Y., Zhang, S.: On the low rank solutions for linear matrix inequalities. *Math. Oper. Res.* **33**(4), 965–975 (2008)
2. Anari, N., Gurvits, L., Gharan, S.O., Saberi, A.: Simply exponential approximation of the permanent of positive semidefinite matrices. In: 2017 IEEE 58th Annual Symposium on Foundations of Computer Science (FOCS), pp. 914–925 (2017)

3. Barvinok, A.: Integration and optimization of multivariate polynomials by restriction onto a random subspace. *Found. Comput. Math.* **7**(2), 229–244 (2007)
4. Egorychev, G.P.: The solution of van der Waerden's problem for permanents. *Adv. Math.* **42**(3), 299–305 (1981)
5. Falikman, D.I.: Proof of the van der Waerden conjecture regarding the permanent of a doubly stochastic matrix. *Math. Notes Acad. Sci. USSR* **29**(6), 475–479 (1981)
6. Gallager, R.G.: Circularly-symmetric gaussian random vectors. <http://www.rle.mit.edu/rgallager/documents/CircSymGauss.pdf>
7. Grier, D., Schaeffer, L.: New hardness results for the permanent using linear optics. In: Proceedings of the 33rd Computational Complexity Conference (San Diego, CA), CCC '18, Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik, June 2018, pp. 1–29
8. Gurvits, L.: Van der Waerden/Schrijver-Valiant like Conjectures and Stable (aka Hyperbolic) Homogeneous Polynomials: One Theorem for all. *Electron. J. Combin.* **15** (2008)
9. Jerrum, M., Sinclair, A., Vigoda, E.: A polynomial-time approximation algorithm for the permanent of a matrix with nonnegative entries. *J. ACM* **51**(4), 671–697 (2004)
10. Keener, R.W.: *Theoretical Statistics: Topics for a Core Course*. Springer Texts in Statistics. Springer, New York (2010)
11. Lobo, M.S., Vandenberghe, L., Boyd, S., Lebret, H.: Applications of second-order cone programming. *Linear Algebra Its Appl.* **284**(1–3), 193–228 (1998)
12. Pappas, A., Révész, S.G.: Linear polarization constants of Hilbert spaces. *J. Math. Anal. Appl.* **300**(1), 129–146 (2004)
13. Pate, T.H.: An inequality involving permanents of certain direct products. *Linear Algebra Its Appl.* **57**, 147–155 (1984)
14. Valiant, L.G.: The complexity of computing the permanent. *Theor. Comput. Sci.* **8**(2), 189–201 (1979)
15. Vandenberghe, L., Boyd, S., Wu, S.-P.: Determinant maximization with linear matrix inequality constraints. *SIAM J. Matrix Anal. Appl.* **19**(2), 499–533 (1998)
16. Zhang, F.: An update on a few permanent conjectures. *Spec. Matrices* **4**(1) (2016)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.